

兔抗 C3 (β chain)多克隆抗体

中文名称: 兔抗 C3 (β chain)多克隆抗体

英文名称: Anti-C3 (β chain) rabbit polyclonal antibody

别 名: ASP; C3a; C3b; AHUS5; ARMD9; CPAMD1

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: C3 (β chain)

反应种属: Human

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

| Background: | Complement component C3 plays a central role in the a ctivation of complement system. Its activation is require d for both classical and alternative complement activation pathways. People with C3 deficiency are susceptible to bacterial infection. Derived from proteolytic degradation of complement C3, C3a anaphylatoxin is a mediator of local inflammatory process. It induces the contraction of smooth muscle, increases vascular permeability and causes histamine release from mast cells and basophilic l |
|---------------|---|
| Applications: | eukocytes. ELISA, WB |



| Name of antibody: | C3 (β chain) |
|------------------------------------|---|
| Immunogen: | Synthetic peptide of human C3 (β chain) |
| Full name: | complement component 3 (β chain) |
| Synonyms: | ASP; C3a; C3b; AHUS5; ARMD9; CPAMD1 |
| SwissProt: | P01024 |
| ELISA Recommended dilution: | 1000-2000 |
| WB Predicted band size: | 70 kDa |
| WB Positive control: | Human serum |
| WB Recommended dilution: | 200-1000 |

