

## 兔抗 CD27 多克隆抗体

中文名称：兔抗 CD27 多克隆抗体

英文名称： Anti-CD27 rabbit polyclonal antibody

别名： T14; S152; Tp55; TNFRSF7; S152. LPFS2

相关类别： 一抗

抗原： CD27

储存： 冷冻（-20℃）

宿主： Rabbit

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

#### Background:

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an import

|                                    |   |
|------------------------------------|---|
|                                    | ant role in the apoptosis induced by this receptor. |
| <b>Applications:</b>               | ELISA, WB, IHC                                      |
| <b>Name of antibody:</b>           | CD27  |
| <b>Immunogen:</b>                  | Fusion protein of human CD27                        |
| <b>Full name:</b>                  | CD27 molecule                                       |
| <b>Synonyms :</b>                  | T14; S152; Tp55; TNFRSF7; S152. LPFS2               |
| <b>SwissProt:</b>                  | P26842  |
| <b>ELISA Recommended dilution:</b> | 2000-5000   |
| <b>IHC positive control:</b>       | Human colon cancer and human thyroid cancer         |
| <b>IHC Recommend dilution:</b>     | 50-200  |
| <b>WB Predicted band size:</b>     | 29 kDa  |
| <b>WB Positive control:</b>        | Raji and lovo cells                                 |
| <b>WB Recommended dilution:</b>    | 500-2000  |



