

CYFIP2 抗原(重组蛋白)

中文名称: CYFIP2 抗原(重组蛋白)

英文名称: CYFIP2 Antigen (Recombinant Protein)

别 名: PIR121

储 存: 冷冻(-20℃)

相关类别: 抗原

概 述:

Fusion protein corresponding to a region derived from C terminal 250 amino acids of human CYFIP2

技术规格:

| Full name: | cytoplasmic FMR1 interacting protein 2 |
|--------------------|---|
| Synonyms: | PIR121 |
| Swissprot: | Q96F07 |
| Gene Accession: | BC011762 |
| Purity: | >85%, as determined by Coomassie blue stained SDS-PAGE |
| Expression system: | Escherichia coli |
| Tags: | His tag C-Terminus, GST tag N-Terminus |
| Background: | CYFIP2 (Cytoplasmic FMR1-interacting protein 2, p53-inducible prote in 121) is a 1,278 amino acid protein encoded by the human gene CYFIP2. CYFIP2 belongs to the CYFIP family and is involved in T-cell adhesion and p53-dependent induction of apoptosis. It interacts with FMR1, FXR1 and FXR2 and is a component of the WAVE1 complex composed of Abi-2, CYFIP2, C3orf10/HSPC300, NAP125 and WASF |



1/WAVE1. Upon binding to activated Rac 1, CYFIP2 causes the complex to dissociate, releasing activated WASF1. The CYFIP2 promoter contains a p53-responsive element that confers p53 binding as well as transcriptional activation of a heterologous reporter. Induced expression of CYFIP2 is sufficient for caspase activation and cellular apoptosis, reminiscent of p53 activation.